

Faculty of Engineering & Technology
First Semester B.E. (C.B.S.) Examination
APPLIED MATHEMATICS—I
Paper—I

Time—Three Hours]

[Maximum Marks—80

INSTRUCTIONS TO CANDIDATES

- (1) All questions carry marks as indicated.
- (2) Use of Non-programmable calculator is permitted.
- (3) Solve Q.No. 1 OR Q.No. 2
Q.No. 3 OR Q.No. 4
Q.No. 5 OR Q.No. 6
Q.No. 7 OR Q.No. 8
Q.No. 9 OR Q.No. 10
Q.No. 11 OR Q.No. 12.

1. (a) If $y = \sin \log (x^2 + 2x + 1)$, prove that :

(i) $(x + 1)^2 y_2 + (x + 1)y_1 + 4y = 0$

(ii) $(x+1)^2 y_{n+2} + (2n+1)(x+1) y_{n+1} + (n^2+4) y_n = 0$ 6

(b) Evaluate $\lim_{x \rightarrow 0} \frac{\sin x^2 - \sin^2 x}{x^4}$ 3

(c) Evaluate $\lim_{x \rightarrow 0} (\cot x)^{\sin x}$ 3

OR

2. (a) Find radius of curvature, centre of curvature and equation to the circle of curvature for the curve $xy(y-x) = 2$ at the point $(1, -1)$. 7

(b) Using Taylor's series, find the value of $\cos 64^\circ$ correct upto four decimal places. 5

3. (a) If $\theta = t^n e^{-t^2/4}$, find what value of n will make $\frac{1}{r^2} \frac{\partial}{\partial t} \left(r^2 \frac{\partial \theta}{\partial t} \right) = \frac{\partial \theta}{\partial t}$. 6

(b) If $u = \tan^{-1} \left(\frac{x^3 + y^3}{x - y} \right)$, then find the value of $x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2}$. 6

(c) If $u = f\left(\frac{x}{y}, \frac{y}{z}, \frac{z}{x}\right)$, find the value of

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z}$$
6

OR

4. (a) Show that the functions :

$$u = \sin^{-1} x + \sin^{-1} y$$

$$\text{and } v = x\sqrt{1-y^2} + y\sqrt{1-x^2}$$

are functionally related. Find the relation between them. 6

(b) Expand $e^x \log(1+y)$ in the neighbourhood of origin by Taylor's series upto six terms of the expansion. 6

(c) Find the volume of the greatest rectangular parallelepiped that can be inscribed in the ellipsoid $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$. 6

5. (a) Find the inverse of matrix by partitioning method :

$$A = \begin{bmatrix} 2 & 3 & 4 \\ 4 & 3 & 1 \\ 1 & 2 & 4 \end{bmatrix}$$
6

(b) Find the rank of matrix :

$$A = \begin{bmatrix} 6 & 1 & 3 & 8 \\ 4 & 2 & 6 & -1 \\ 10 & 3 & 9 & 7 \\ 16 & 4 & 12 & 15 \end{bmatrix}$$

6

OR

6. (a) Investigate the values of λ & μ so that the system of equations :

$$2x + 3y + 5z = 9$$

$$7x + 3y - 2z = 8$$

$$2x + 3y - \lambda z = \mu$$
 have

(i) No solution

(ii) Unique solution

(iii) Infinite solutions.

6

(b) Solve the system of equations :

$$x + y + z = 3$$

$$x + 2y + 3z = 4$$

$$x + 4y + 9z = 6$$

by adjoint method.

6

MMW-9682

4

(Contd.)

7. (a) Solve $\frac{dy}{dx} - \frac{\tan y}{1+x} = (1+x)e^x \sec y$. 4

(b) Solve $(1+y^2) dx = (\tan^{-1}y - x) dy$. 4

(c) Solve :

$$(y^2 e^{xy} + 4x^3) dx + (2xy e^{xy} - 3y^2) dy = 0. 4$$

OR

8. (a) Solve $xy^2 (p^2 + 2) = 2py^3 + x^3$. 3

(b) Solve $y = 2px + p^4 x^2$.

(c) A 20 ohm resistor is connected in series with capacitor of 0.01 farad and emf E volt given by $40e^{-3t} + 20e^{-4t}$. If $q = 0$ at $t = 0$, show that the maximum charge on the capacitor is 0.25 coulomb.

9. (a) Solve $\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + y = e^x \cdot \cos 2x$.

(b) Solve $(D^2 + 2D + 1)y = 4e^{-x} \log x$ by method of variation of parameters. 6

(c) $x^2 \frac{d^2y}{dx^2} - 2x \frac{dy}{dx} - 4y = x^2 + 2 \log x$. 6

OR

MMW-9682

5

(Contd.)

10. (a) An emf $E \sin pt$ is applied at $t = 0$ to a circuit containing a capacitance C & inductance L . The current i satisfies the equation

$$L \frac{di}{dt} + \frac{1}{C} \int i dt = E \sin pt$$

If $p^2 = \frac{1}{LC}$ and initially the current i and the charge q be zero, show that the current at time t is $\frac{Et}{2L} \sin pt$, where $i = \frac{dq}{dt}$. 7

- (b) A mechanical system with two degree of freedom satisfies the equations :

$$2 \frac{d^2x}{dt^2} + 3 \frac{dy}{dt} = 4, \quad 2 \frac{d^2y}{dt^2} - 3 \frac{dx}{dt} = 0.$$

Obtain expressions for x and y in terms of t , given $x, y, \frac{dx}{dt}, \frac{dy}{dt}$ all vanish at $t = 0$. 6

- (c) Solve $\frac{d^2y}{dx^2} = \sec^2 y \tan y$, given that $y = 0$ and

$$\frac{dy}{dx} = 1, \text{ when } x = 0. \quad 5$$

11. (a) Using De-Moivre's theorem, solve :

$$x^5 + x^4 + x^3 + x^2 + x + 1 = 0. \quad 4$$

- (b) If $\sin(A + iB) = x + iy$, then prove that :

$$(i) \frac{x^2}{\cosh^2 B} + \frac{y^2}{\sinh^2 B} = 1$$

$$(ii) \frac{x^2}{\sin^2 A} - \frac{y^2}{\cos^2 A} = 1.$$

OR

12. (a) Prove that :

$$\cos^6 \theta + \sin^6 \theta = \frac{1}{8}(3 \cos 4\theta + 5).$$

- (b) Find the modulus and argument of $(1 + i)^{1-i}$.