

NKT/KS/17/5199

Bachelor of Science (B.Sc.) Semester—VI (C.B.S.) Examination
M₁₂—DISCRETE MATHEMATICS AND ELEMENTARY NUMBER THEORY
Paper-2

(Mathematics)

Time : Three Hours]

[Maximum Marks : 60

N.B. :— (1) Solve all the **FIVE** questions.

(2) All questions carry equal marks.

(3) Question Nos. 1 to 4 have an alternative. Solve each question in full or its alternative in full.

UNIT-I1. (A) Given that $A = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$ and a relation R on A is such that $R = \{(x, y) / x + y = 10\}$. Show that R is neither reflexive nor transitive but it is symmetric.

6

(B) Let (L, \leq) be a Lattice. For any $a, b, c, \in L$; if $*$ and \oplus are operations of meet and join, then prove that :

$$b \leq c \Rightarrow \begin{cases} a * b & \leq & a * c \\ a \oplus b & \leq & a \oplus c \end{cases}$$

6


OR

(C) Define a distributive lattice and show that every chain is a distributive lattice.

6

(D) Define : Indegree, Outdegree and Total degree of node. Find all the indegree and outdegree of the digraph.

6



UNIT-II

2. (A) Prove that if g is greatest common divisor of b and c , then there exist integers x_0 and y_0 such that $g = (b, c) = bx_0 + cy_0$. 6
- (B) Find the value of x and y that satisfy the equation :
 $243x + 198y = 9$ 6

OR

- (C) Let p be an odd prime and $(a, p) = 1$. Then prove that
 $a^{p-1} \equiv 1 \pmod{p}$. 6
- (D) Evaluate :
 $(n, n+1)$ and $[n, n+1]$. 6

UNIT-III

3. (A) Define Legendre's Symbol. Prove that if p is an odd prime, then

$$\left(\frac{-1}{p}\right) = \begin{cases} 1 & \text{if } p \equiv 1 \pmod{4} \\ -1 & \text{if } p \equiv 3 \pmod{4} \end{cases}$$
 6

- (B) Evaluate $\left(\frac{-42}{61}\right)$ and $\left(\frac{30}{71}\right)$. 6

OR

- (C) Solve the congruence if it is solvable :
 $x^2 \equiv 7 \pmod{31}$. 6
- (D) Prove that the quadratic congruence :
 $x^2 \equiv a \pmod{p}$ where p is odd positive prime integer with $a \not\equiv kp$ for any integer k , has exactly two solutions, provided the solution exist. 6

UNIT-IV

4. (A) Prove that the positive primitive solution of $x^2 + y^2 = z^2$ with y even are given by $x = r^2 - s^2$,
 $y = 2rs$, $z = r^2 + s^2$
 where r, s are arbitrary positive integers of opposite parity and $r > s$ and $(r, s) = 1$. 6

- (B) Find all the primitive Pythagorean triples x, y, z such that $z - y = 1$. 6

OR

- (C) Construct Farey Sequence F_6 , given that $F_1 = \left\{ \frac{0}{1}, \frac{1}{1} \right\}$. 6

- (D) If $\frac{a}{b}$ and $\frac{a'}{b'}$ are two consecutive terms in F_n with $\frac{a}{b}$ to the left of $\frac{a'}{b'}$, then prove $a'b - ab' = 1$. 6

QUESTION-V

5. (A) Define an equivalence relation. Give an example of equivalence relation. $1\frac{1}{2}$
- (B) Define simple graph and multi-graph. $1\frac{1}{2}$
- (C) Prove that $4x \mid (n^2 + 2)$ for any integer n . $1\frac{1}{2}$
- (D) Define Reduced Residue System. Find reduced residue system for $m = 12$. $1\frac{1}{2}$
- (E) Prove that $\left(\frac{a}{p} \right) = 1$. $1\frac{1}{2}$
- (F) Define greatest integer function $[x]$. Find $[-3.5]$ and $[12]$. $1\frac{1}{2}$
- (G) Test the solvability of $4x + 8y = 2$. $1\frac{1}{2}$
- (H) Prove that terms in a Farey sequence are in monotonically increasing order. $1\frac{1}{2}$

NKT/KS/17/5200

Bachelor of Science (B.Sc.) Semester—VI (C.B.S.) Examination**M₁₂—DIFFERENTIAL GEOMETRY****Paper-2****(Mathematics)**

Time : Three Hours]

[Maximum Marks : 60

N.B. :— (1) Solve all the **FIVE** questions.

(2) All questions carry equal marks.

(3) Question Nos. 1 to 4 have an alternative. Solve each question in full or its alternative in full.

UNIT-I1. (A) Prove that Darboux vector \vec{d} is constant if K and τ are constant and the d has a fixed directionin $\frac{K}{\tau}$ is constant.

6

(B) Prove that the Frenet-Serret formulae :

(i) $\frac{d\vec{t}}{ds} = K\vec{n}$

(ii) $\frac{d\vec{n}}{ds} = -\tau\vec{b} - K\vec{t}$

(iii) $\frac{d\vec{b}}{ds} = \tau\vec{n}$

where K and τ are curvature and torsion of the space curve.

6

OR

(C) Show that the tangent to the locus of the centres of the osculating sphere passes through the centre of the osculating circle.

6

(D) Prove that the necessary and sufficient condition that a curve to be a helix is that ratio of curvature to torsion is constant at all points.

6

UNIT-II

2. (A) Find the involutes and evolutes of the circular helix :

$$\vec{r} = (a \cos \theta, a \sin \theta, b\theta). \quad 6$$

- (B) Show that the curvature K_1 and torsion τ_1 of an involute \tilde{c} of c are given by

$$K_1^2 = \frac{\tau^2 + K^2}{K^2 (c-s)^2}, \quad \tau_1 = \frac{K\tau' - K'\tau}{K(c-s)(K^2 + \tau^2)}$$

where K and τ are the curvature and the torsion of the curve. 6

OR

- (C) Find the principal curvatures at the origin of the paraboloid

$$2\zeta = 5x^2 + 4xy + 2y^2.$$

Also find the curvature of the section $y = x$. 6

- (D) If $\dot{\alpha}\dot{\beta} - \dot{\beta}\dot{\alpha} \neq 0$, show that the line $x = a\zeta + \alpha$, $y = b\zeta + \beta$ generates a skew ruled surface and find its equation where a, b and α, β are functions of u . 6

UNIT-III

3. (A) Determine the unit normals and the fundamental forms of the surface

$$\vec{r} = (a \cos u, a \sin u, bv). \quad 6$$

- (B) On the paraboloid $x^2 - y^2 = \zeta$, find the orthogonal trajectories of the section by the planes $z = \text{constant}$. 6

OR

- (C) Obtain Gauss's formulae for \vec{r}_{11} , \vec{r}_{12} , \vec{r}_{22} , where \vec{r} is the position vector of any point of a surface and suffixes 1 and 2 denotes differentiation with regard to u and v respectively. 6

- (D) Prove that the Rodrigue's formula $Kd\vec{r} + d\vec{N} = 0$

where K is the principle curvature. 6

UNIT-IV

4. (A) Define geodesic and further obtain the differential equation of the geodesic. 6
- (B) Prove that the curves of the family $\frac{v^3}{u^2} = \text{constant}$ are geodesics on a surface with the metric.
 $v^2 du^2 - 2 uv du dv + 2u^2 dv^2, u > 0, v > 0.$ 6

OR

- (C) Find the Gaussian curvature at a point (u, v) of the anchor ring,
 $r = ((b + a \cos u) \cos v, (b + a \cos u) \sin v, a \sin u), 0 < u, v < 2\pi.$ 6
- (D) Prove Bonnet's theorem for a curve on a surface that $w' + \tau = \tau_g$, where w is the normal angle and τ_g is the torsion of the geodesic tangent. 6

QUESTION-V

5. (A) Define Osculating Plane. 1½
- (B) Define Fundamental plane at a point P whose position vector is \vec{r} on the space curve \vec{R} . 1½
- (C) Define Involute and Evolute. 1½
- (D) Define Developable Surface. 1½
- (E) Define a third fundamental form. 1½
- (F) State Euler's theorem on normal curvature. 1½
- (G) State Gauss-Bonnet theorem. 1½
- (H) Define geodesic polar coordinates for the geodesic metric $ds^2 = du^2 + G(u, v) dv^2.$ 1½

NKT/KS/17/5201

Bachelor of Science (B.Sc.) Semester—VI (C.B.S.) Examination**M₁₂—SPECIAL THEORY OF RELATIVITY****Paper–2****(Mathematics)**

Time : Three Hours]

[Maximum Marks : 60

N.B. :— (1) Solve all the **FIVE** questions.

(2) All questions carry equal marks.

(3) Question Nos. 1 to 4 have an alternative. Solve each question in full or its alternative in full.

UNIT–I

1. (A) What is an inertial frame ? Show that in an inertial frame, a body not under influence of any forces, moves in a straight line with constant velocity. 6

- (B) Prove that $\nabla^2 - \frac{1}{C^2} \frac{\partial^2}{\partial t^2}$ is invariant under special Lorentz transformations. 6

OR

- (C) Explain Lorentz-Fitzgerald contraction idea. How was this idea used to account for negative result of the Michelson-Morley experiment ? 6

- (D) Assume that the Lorentz transformations are the linear transformations of the form $x' = Ax + Bt$ and $t' = Dx + Et$, $E > 0$. Considering $x^2 - c^2 t^2 = x'^2 - c^2 t'^2$ and the motion of the origin of S, deduce the Lorentz transformations. 6

UNIT–II

2. (A) Obtain the transformation equations for the components of particle acceleration by using special Lorentz transformations. 6

- (B) Explain the phenomenon of time dilation in special theory of relativity. 6

OR

- (C) Obtain the transformation of Lorentz contraction factor $\left(1 - \frac{u^2}{c^2}\right)^{\frac{1}{2}}$ in the two inertial frames of references. 6
- (D) Prove that simultaneity is relative but not an absolute in special relativity. 6

UNIT-III

3. (A) Define symmetric and skew symmetric covariant tensors of order two. Show that any tensor of the second order may be expressed as the sum of a symmetric tensor and skew symmetric tensor. 6
- (B) Define Kronckar Delta ∂_s^r . Show that ∂_s^r is a mixed tensor of order two. 6

OR

- (C) Define space-like interval. Prove that there exists an inertial system S' in which the two events occur at one and the same time if the interval between two events is space-like. 6
- (D) Define four tensor. Obtain the transformations of the components of a symmetrical four tensor T^{11} under the Lorentz transformations. 6

UNIT-IV

4. (A) If m is the relativistic mass of a particle moving with velocity u relative to an inertial frame s, then prove that :

$$m = \frac{m_0}{\left(1 - \frac{u^2}{c^2}\right)^{\frac{1}{2}}}$$

where m_0 is the rest mass of the particle. 6

- (B) Define four-velocity. Show that the four velocity, in component form, can be expressed as :

$$u^i = \left(\frac{\bar{u}}{c \sqrt{1 - u^2/c^2}}, \frac{1}{\sqrt{1 - u^2/c^2}} \right)$$

where $\bar{u} = (u_x, u_y, u_z)$. 6

OR

- (C) Derive the wave equation for the propagation of the electric field strength \vec{E} and the magnetic field strength \vec{H} in free space with velocity of light. 6
- (D) Obtain the transformation equations of the electromagnetic four-potential vector. 6

QUESTION-V

5. (A) Discuss the outcome of Michelson-Morley experiment regarding fringe shift and stationary ether. 1½
- (B) Show that the circle $x'^2 + y'^2 = a^2$ in the frame of reference s' is measured to be an ellipse in s if s' moves with uniform velocity relative to s . 1½
- (C) Derive Einstein's velocity addition law. 1½
- (D) An electron is moving with a speed of $0.85c$ in a direction opposite to that of a moving photon. Calculate relative velocity. 1½
- (E) Define :
Free suffix and dummy suffix in a tensor quantity. 1½
- (F) Show that two events which are separated by a time-like interval cannot occur simultaneously in any inertial system. 1½
- (G) Show that the four velocity of a particle is a unit time-like vector. 1½
- (H) Prove that the energy momentum tensor T^j_i is symmetric. 1½